

### Summary diagram for medical examinations, treatments and activity resumption

	0 to 3 Months	3 to 6 Months	7 to 12 Months	18 months and +	2 years	3 years	4 years and +
Check-up by the paediatrician or gastroenterologist (at least)	1 x week	1x every 2 weeks	1x Month	All 3 Months			
Check-up at the HUG (at least)	X	X	X	X	X	X	X

	3 Months	6 Months	9 Months	1 year	18 months	2 years	3 years	4 and +
<b>immunosuppressor °•</b>	Dose according to protocol, treatment according to situation			Lifetime treatment				
<b>Prednisone ®</b>	According to diagram		More if indication					
<b>Nexium / oméprazole ®</b>	With Prédnisone							
<b>Valcyte ®</b>	6 Months							
<b>Nopil ®</b>				1 year				
<b>Aspirine ®</b>	3 Months		More if surgical indication					
<b>Ursofalk ®</b>				1 year		More if indication		
<b>Other treatments</b>	amlodipine, spironolactone, vitamins, iron, magnesium, calcium and all other treatments will be prescribed as needed							

°• Tacrolimus, everolimus, MMF, ciclosporine

	3 Months	6 Months	9 Months	1 year	18 months	2 years	3 years	4 and +
<b>Nursery</b>		* 1						
<b>School</b>	* 1							
<b>Sport</b>	* 2	no restriction (except swimming pool)						
<b>Sea, private swimming pool</b>								
<b>Public swimming pool, spa / wellness, lake or river</b>								
* 1	Depending on the situation							
* 2	6 weeks without intensive or contact sports, 3 months without trampoline or trottiers							